

A Geometric Construction of Getzler's Elliptic Relation

Rahul Pandharipande

12 May 1997

0 Introduction

Let $\overline{M}_{1,4}$ be the moduli space of Deligne-Mumford stable 4-pointed elliptic curves. E. Getzler has determined the natural $\mathbb{S}_4$ -module structure on the vector space $H^*(\overline{M}_{1,4}, \mathbb{Q})$ using modular operads and Deligne's mixed Hodge theory [G1]. The dimension of the $\mathbb{S}_4$ -invariant space $H^4(\overline{M}_{1,4}, \mathbb{Q})^{\mathbb{S}_4}$ is 7. $\overline{M}_{1,4}$ has a natural stratification by dual graph type. Define an $\mathbb{S}_4$ -invariant stratum of $\overline{M}_{1,4}$ to be an $\mathbb{S}_4$ -orbit of closed strata of $\overline{M}_{1,4}$. The number of invariant dimension 2 strata in $\overline{M}_{1,4}$ is 9. The classes of these invariant strata in cohomology must therefore satisfy at least 2 linear relations. The first relation is evident. Let Δ_0 be the boundary stratum with generic element corresponding to a 4-pointed nodal rational curve. There is a natural map $\overline{M}_{0,6} \rightarrow \Delta_0$ obtained by identifying the markings 5 and 6. Pushing forward the basic divisor linear equivalence on $\overline{M}_{0,6}$ to Δ_0 yields a relation among the dimension 2 boundary strata of $\overline{M}_{1,4}$ contained in Δ_0 . In [G2], Getzler computes the 9×9 intersection pairing of the invariant strata in $\overline{M}_{1,4}$. This intersection matrix is found to have rank 7. The invariant strata therefore span $H^4(\overline{M}_{1,4}, \mathbb{Q})^{\mathbb{S}_4}$. The null space of the intersection matrix is computed to find a new relation among these strata in cohomology.

A direct construction of Getzler's new relation via a rational equivalence in the Chow group $A_2(\overline{M}_{1,4}, \mathbb{Q})$ is presented here. The idea is to equate cycles corresponding to different degenerations of elliptic curves in the Chow group of a space of admissible covers. The push-forward of this equivalence to $\overline{M}_{1,4}$ yields Getzler's equation.

The strategy of the construction is the following. First, select a point:

$$\zeta = [\mathbf{P}^1, p_1, p_2, p_3, p_4] \in M_{0,4}.$$

Consider the space of degree 2 maps $\pi : E \rightarrow \mathbf{P}^1$ from 4-pointed nonsingular elliptic curves to $\mathbf{P}^1$ satisfying:

- (i) π is not ramified at the markings of E ,
- (ii) The i^{th} marking of E lies over p_i .

This space has dimension 4 and may be compactified (after ordering the ramification points) by the space of pointed admissible double covers $\overline{H}_\zeta$. These spaces are discussed in Section 1. There are two natural dominant maps:

$$\begin{aligned}\lambda : \overline{H}_\zeta &\rightarrow \overline{M}_{1,4}, \\ \rho : \overline{H}_\zeta &\rightarrow \text{Sym}^4(\mathbf{P}^1).\end{aligned}$$

The map λ is the map to moduli while $\rho([\pi])$ is defined by the degree 4 ramification divisor of π . There are two natural surfaces in $\text{Sym}^4(\mathbf{P}^1)$: the loci where the generic divisor shapes are 2+2 and 1+3 respectively. These surfaces are linearly equivalent (up to a scalar) in $A_2(\text{Sym}^4(\mathbf{P}^1), \mathbb{Q})$ since $A_2(\text{Sym}^4(\mathbf{P}^1), \mathbb{Q}) = \mathbb{Q}$. The intersection pull-back via ρ of these surfaces yields a rational equivalence of codimension 2 cycles in $\overline{H}_\zeta$. This is a *boundary* linear equivalence because a double cover by a nonsingular elliptic curve is branched over 4 distinct points. The λ push-forward of this relation yields Getzler's relation in $A_2(\overline{M}_{1,4}, \mathbb{Q})$.

C. Faber has determined the Chow ring of $\overline{M}_3$ in [Fa]. There are 12 dimension 2 boundary strata. In [Fa], the rank $A_2(\overline{M}_3, \mathbb{Q}) = 7$ is computed and the strata are shown to span. Four geometric relations are constructed among the dimension 2 boundary strata. Faber deduces a fifth relation via the ring axioms and further knowledge of $A_*(\overline{M}_3, \mathbb{Q})$. There is a map $\tau : \overline{M}_{1,4} \rightarrow \overline{M}_3$ obtained by identifying the marked point 1 with 2 and 3 with 4. The τ push-forward of Getzler's relation is easily seen to yield a relation among the dimension 2 strata of $\overline{M}_3$. This provided a geometric construction of Faber's fifth relation.

Getzler's new relation yields a differential equation for elliptic Gromov-Witten invariants analogous to the WDVV-equations in genus 0 (see [G2]). Eguchi, Hori, and Xiong and, independently, S. Katz have proposed the full (gravitational) potential function is annihilated by a certain representation of the Virasoro algebra. The Virasoro conjecture (together with the topological recursion relations in genus 1) predicts a remarkably simple recursion for the elliptic Gromov-Witten invariants of $\mathbf{P}^2$ [EHX]. In the last section, this elliptic recursion is proven via Getzler's equation and the WDVV-equations.

The author wishes to thank D. Abramovich, P. Belorousski, C. Faber, E. Getzler, T. Graber, S. Katz, and A. Kresch for conversations about $\overline{M}_{1,4}$ and related issues. This research was completed at the Mittag-Leffler Institute. The author was partially supported by an NSF post-doctoral fellowship.

1 Moduli of maps

An *admissible cover of degree 2 with 4 marked points and 4 branch points* consists of a morphism $\pi : E \rightarrow D$ of pointed curves

$$[E, P_1, \dots, P_4], [D, p_1, \dots, p_4, q_a, \dots, q_d]$$

satisfying the following conditions:

- (i) E is a connected, reduced, nodal curve of arithmetic genus 1.

- (ii) The markings P_i lie in the nonsingular locus E_{ns} .
- (iii) $\pi(P_i) = p_i$.
- (iv) $[D, p_1, \dots, p_4, q_a, \dots, q_d]$ is an 8-pointed stable curve of genus 0.
- (v) $\pi^{-1}(D_{ns}) = E_{ns}$.
- (vi) $\pi^{-1}(D_{sing}) = E_{sing}$.
- (vii) $\pi|_{E_{ns}}$ is étale except over the points q_j where π is simply ramified.
- (viii) If $x \in E_{sing}$, then:
 - (a) $x \in E_1 \cap E_2$ where E_1, E_2 are distinct components of E .
 - (b) $\pi(E_1), \pi(E_2)$ are distinct components of D .
 - (c) The ramification numbers at x of the two morphisms:

$$\pi : E_1 \rightarrow \pi(E_1), \quad \pi : E_2 \rightarrow \pi(E_2)$$

are equal.

Let $\overline{H}$ be the space of 4-pointed admissible degree 2 covers of $\mathbf{P}^1$ branched at 4 points.

$\overline{H}$ is an irreducible variety parametrizing admissible covers. There are natural maps

$$\lambda : \overline{H} \rightarrow \overline{M}_{1,4}, \quad \pi : \overline{H} \rightarrow \overline{M}_{0,8} \tag{1}$$

obtained from the domain and range of the admissible cover respectively. The projection π is a finite map to $\overline{M}_{0,8}$. For each marking $i \in \{1, 2, 3, 4\}$, there is a natural $\mathbb{Z}/2\mathbb{Z}$ -action given by switching the sheet of the i^{th} -marking of E . These actions induce a product action in which the diagonal $\triangle$ acts trivially. Define the group $\mathbf{G}$ by:

$$\mathbf{G} = (\mathbb{Z}/2\mathbb{Z})^4 / \triangle.$$

The action of $\mathbf{G}$ on $\overline{H}$ is generically free and commutes with the projection π . Therefore, the quotient $\overline{H}/\mathbf{G}$ naturally maps to $\overline{M}_{0,8}$. In fact, since $\overline{H}/\mathbf{G} \rightarrow \overline{M}_{0,8}$ is finite and birational, it is an isomorphism.

Spaces of admissible covers were defined in [HM]. The methods there may be used to construct spaces of pointed admissible covers. A quick route to $\overline{H}$ via Kontsevich's space of stable maps will be presented here.

Let $\mu : U \rightarrow \overline{M}_{0,8}$ be the 8-pointed universal curve. Let $\overline{M}_{1,4}(\mu, 2)$ be the relative space of stable maps of double covers of the universal curve. $\overline{M}_{1,4}(\mu, 2)$ is a projective variety (see [K], [KM], [FP], [BM]). Let H denote the quasi-projective subvariety of $\overline{M}_{1,4}(\mu, 2)$ corresponding to 4-pointed maps $\mu : E \rightarrow D$ satisfying:

- (i) E is a nonsingular 4-pointed elliptic curve, D is a nonsingular 8-pointed $\mathbf{P}^1$.

- (ii) The markings of E lie over the first four markings of D .
- (iii) The four ramification points of μ lie over the last four markings of D .

H is an étale 8-sheeted cover of $M_{0,8}$ with a free $\mathbf{G}$ -action. Let $\overline{H}$ be the closure of H in $\overline{M}_{1,4}(\mu, 2)$. The points of the closure are seen to parameterize pointed admissible covers. The basic maps (1) are obtained from the restriction of maps defined on $\overline{M}_{1,4}(\mu, 2)$. The $\mathbf{G}$ -action is deduced from the universal properties of $\overline{M}_{1,4}(\mu, 2)$. $\overline{H}$ and $\overline{M}_{0,8}$ are 5 dimensional spaces.

Let $\zeta = [\mathbf{P}^1, p_1, p_2, p_3, p_4] \in M_{0,4}$. Let X_ζ be the fiber over ζ of the natural map $\overline{M}_{0,8} \rightarrow \overline{M}_{0,4}$ determined by the first four markings. X_ζ is a nonsingular 4 dimensional variety. Another way to view X_ζ is the following. Let $\mathbf{P}^1[8]$ be the Fulton-MacPherson configuration space [FM]. Let $\gamma_1, \dots, \gamma_4$ be the first four projections $\gamma_i : \mathbf{P}^1[8] \rightarrow \mathbf{P}^1$. X_ζ is naturally identified with the intersection:

$$\gamma_1^{-1}(p_1) \cap \gamma_2^{-1}(p_2) \cap \gamma_3^{-1}(p_3) \cap \gamma_4^{-1}(p_4) \subset \mathbf{P}^1[8].$$

In particular, the four markings corresponding to the ramification points yield four natural projections $\gamma_a, \dots, \gamma_d$ from X_ζ to $\mathbf{P}^1$.

X_ζ is a compactification of the space of 8 points on $\mathbf{P}^1$ where the first four have a specified cross ratio. The boundary of X_ζ is a divisor with normal crossings. Moreover, X_ζ has a stratification by graph type.

Let $\overline{H}_\zeta = \pi^{-1}(X_\zeta)$. $\overline{H}_\zeta$ is a natural compactification of the space of 4-pointed nonsingular elliptic double covers of $\mathbf{P}^1$ satisfying:

- (i) The ramification points on E are ordered and distinct from the markings.
- (ii) The markings lie over a fixed 4-tuple $\zeta \in M_{0,4}$.

$\mathbf{G}$ acts on $\overline{H}_\zeta$ with quotient $\overline{H}_\zeta/\mathbf{G} = X_\zeta$. The restriction $\lambda : \overline{H}_\zeta \rightarrow \overline{M}_{1,4}$ is a generically finite dominant map of degree 48.

2 Two Surfaces

Let $S_{2,2}$ and $S_{1,3}$ be the irreducible surfaces in $Sym^4(\mathbf{P}^1) \cong \mathbf{P}^4$ corresponding to divisors of shape $2 + 2$ and $1 + 3$ respectively. The degrees of these surfaces are 4 and 6 respectively. Therefore,

$$\frac{[S_{2,2}]}{4} = \frac{[S_{1,3}]}{6}$$

in $A_2(Sym^4(\mathbf{P}^1))$. Chow groups here will always be taken with $\mathbb{Q}$ -coefficients.

Index four $\mathbf{P}^1$'s by the letters $\{a, b, c, d\}$. Define:

$$\mathbf{P}_4^1 = \mathbf{P}_a^1 \times \dots \times \mathbf{P}_b^1.$$

The following convenient notation will be used for the diagonal subvarieties of $\mathbf{P}_4^1$. Let (ab) denote diagonal where the coordinates of a and b coincide.

Let $(ab, cd) = (ab) \cap (cd)$. Similarly, let (abc) denote the diagonal where the coordinates of a , b , and c coincide. Let $\mathbf{P}_4^1 \rightarrow \text{Sym}^4(\mathbf{P}^1)$ be the $\mathbb{S}_4$ -quotient map. The inverse image of $S_{2,2}$ in $\mathbf{P}_4^1$ is $(ab, cd) \cup (ac, bd) \cup (ad, bc)$. Similarly, the inverse image of $S_{1,3}$ is $(abc) \cup (acd) \cup (abd) \cup (bcd)$.

Lemma 1 *Let $\mathbf{A}$ be a finite group. Let Z be an irreducible algebraic variety with an $\mathbf{A}$ -action and a quotient morphism $\alpha : Z \rightarrow Z/\mathbf{A}$. There exists a pull-back $\alpha^* : A_*(Z/\mathbf{A}) \rightarrow A_*(Z)$ defined by:*

$$\alpha^*([V]) = |\text{Stab}(V)| \cdot [\alpha^{-1}(V)_{\text{red}}]$$

where V is an irreducible subvariety of $Z/\mathbf{A}$, the scheme $\alpha^{-1}(V)_{\text{red}}$ is the reduced preimage of V , and $|\text{Stab}(V)|$ is the size of the generic stabilizer of points over V .

By Lemma 1, there is an equality in $A_2(\mathbf{P}_4^1)$:

$$[ab, cd] + [ac, bd] + [ad, bc] = [abc] + [acd] + [abd] + [bcd]. \quad (2)$$

This equality can of course be checked directly. Lemma 1 will be used in Section 4. See [V] for a proof of Lemma 1.

3 Refined Intersection

The notation of Sections 1 and 2 is followed here. The four projections $\gamma_a, \dots, \gamma_d$ from X_ζ to $\mathbf{P}^1$ yield a map:

$$\Gamma : X_\zeta \rightarrow \mathbf{P}_4^1.$$

The pull-back of the relation (2) by the map $\Gamma^* : A^*(\mathbf{P}_4^1) \rightarrow A^*(X_\zeta)$ is now analyzed.

First, the left side (2) is considered. The scheme theoretic inverse $\Gamma^{-1}(ab)$ is a reduced union of boundary divisors of X_ζ . Therefore, since the boundary in X_ζ has normal crossings,

$$\Gamma^{-1}(ab, cd) = \Gamma^{-1}(ab) \cap \Gamma^{-1}(cd)$$

is a reduced union of 2 and 3 dimensional strata X_ζ . In fact, in the notation of Appendices B and C ,

$$\Gamma^{-1}(ab, cd) = S_1 \cup S_2 \cup S_2 \cup S_3 \cup T_3 \cup T_4.$$

S_2 occurs with 2 different round marking assignments. While the strata S_i are of expected dimension 2, T_3 and T_4 are of excess dimension. Excess intersection theory yields a dimension 2 cycle supported on T_3 and T_4 . There is a bundle sequence on the divisor $T = T_3 \cup T_4$ which is exact away from the 1 dimensional intersections of T with the 2 dimensional components of $\Gamma^{-1}(ab, cd)$:

$$0 \rightarrow \mathcal{O}_T(T) \rightarrow \Gamma^*(N_{ab, cd}) \rightarrow L \rightarrow 0.$$

Here, $N_{ab,cd}$ is the normal bundle of (ab, cd) in $\mathbf{P}_4^1$, the first map is obtained from the differential of Γ , and L is the excess bundle. The first Chern class of L is the excess cycle on T (see [Fu]). A standard calculation yields the excess cycle:

$$2R_1 + \frac{1}{2}R_2 + 2R_3 + 2R_4 + S_4 + S_7 + S_8.$$

The intersection pull-back is thus determined by:

$$\Gamma^*[ab, cd] = 2R_1 + \frac{1}{2}R_2 + 2R_3 + 2R_4 + S_1 + 2S_2 + S_3 + S_4 + S_7 + S_8.$$

The above equalities are in $A_2(X_\zeta)/K$ (see Appendix B).

The scheme theoretic inverse $\Gamma^{-1}(abc)$ is a union of boundary divisors of X_ζ :

$$\Gamma^{-1}(abc) = T_1 \cup T_2 \cup T_3 \cup T_4.$$

The excess bundle L is again determined by the natural exact sequence on $T = \Gamma^{-1}(abc)$:

$$0 \rightarrow \mathcal{O}_T(T) \rightarrow \Gamma^*(N_{abc}) \rightarrow L \rightarrow 0.$$

A calculation yields the excess class in $A_2(X_\zeta)/K$:

$$\begin{aligned} \Gamma^*[abc] &= \frac{6}{10}R_1 + \frac{1}{2}R_2 + \frac{13}{10}R_3 + \frac{1}{2}R_4 + \frac{6}{10}R_5 + \frac{1}{10}R_6 + \frac{1}{6}R_7 \\ &\quad + S_4 + S_5 + S_6 + S_7 + S_8 + S_9. \end{aligned}$$

The Γ pull-back of the left side of (2) in $A_2(X_\zeta)/K$ is simply $3\Gamma^*[ab, cd]$. Similarly, the pull-back of the right side is $4\Gamma^*[abc]$. Hence, the above calculation yield an equality in $A_2(X_\zeta)/K$:

$$\begin{aligned} \frac{36}{10}R_1 - \frac{1}{2}R_2 + \frac{8}{10}R_3 + 4R_4 - \frac{24}{10}R_5 - \frac{4}{10}R_6 - \frac{4}{6}R_7 = \quad (3) \\ -3S_1 - 6S_2 - 3S_3 + S_4 + 4S_5 + 4S_6 + S_7 + S_8 + 4S_9. \end{aligned}$$

4 Push-Forwards

The application of $\lambda_*\pi^* : A_2(X_\zeta) \rightarrow A_2(\overline{M}_{1,4})$ to relation (3) yields a boundary relation in $A_2(\overline{M}_{1,4})$. The calculation of π^* is obtained by Lemma 1. It is checked the generic stabilizers of $\mathbf{G}$ over the strata are trivial except in the following cases:

$$\begin{aligned} |\text{Stab}(R_4)| &= 2 \\ |\text{Stab}(R_5)| &= 2 \\ |\text{Stab}(R_6)| &= 2 \\ |\text{Stab}(R_7)| &= 4 \\ |\text{Stab}(S_6)| &= 2 \\ |\text{Stab}(S_9)| &= 2. \end{aligned}$$

The elements $\lambda_*\pi^*(R_i)$ are tabulated below. See Appendix A for the notation in $A_2(\overline{M}_{1,4})$. These elements lie in the linear span of the 4 invariant strata of $A_2(\overline{M}_{1,4})$ with nonsingular elliptic components.

$$\begin{aligned}
\lambda_*\pi^*(R_1) &= 2\Delta_{2,4} + 6\Delta_{3,4} \\
\lambda_*\pi^*(R_2) &= 96\Delta_{2,2} + 32\Delta_{2,3} \\
\lambda_*\pi^*(R_3) &= 12\Delta_{2,4} \\
\lambda_*\pi^*(R_4) &= 16\Delta_{2,3} \\
\lambda_*\pi^*(R_5) &= 12\Delta_{3,4} \\
\lambda_*\pi^*(R_6) &= 32\Delta_{2,4} + 12\Delta_{3,4} \\
\lambda_*\pi^*(R_7) &= 48\Delta_{2,3}
\end{aligned}$$


The elements $\lambda_*\pi^*(S_i)$ lie in the linear span of the 5 invariant strata contained in the boundary divisor $\Delta_0 \subset \overline{M}_{1,4}$ corresponding to a 4-pointed nodal rational curve.

$$\begin{aligned}
\lambda_*\pi^*(S_1) &= \frac{2}{3}\Delta_{0,2} + 2\Delta_a + \frac{16}{3}\Delta_b \\
\lambda_*\pi^*(S_4) &= 4\Delta_{0,2} \\
\lambda_*\pi^*(S_5) &= 2\Delta_{0,4} \\
\lambda_*\pi^*(S_6) &= 2\Delta_{0,3} \\
\lambda_*\pi^*(S_7) &= 2\Delta_{0,3}
\end{aligned}$$

The push-forwards $\lambda_*\pi^*(S_2)$, $\lambda_*\pi^*(S_3)$, $\lambda_*\pi^*(S_8)$, and $\lambda_*\pi^*(S_9)$ all vanish.

In the above formulas, the ordinary coarse moduli fundamental classes in $\overline{M}_{1,4}$ are used on the right. The orbifold classes differ by a factor of two for $\Delta_{2,4}$, $\Delta_{3,4}$, and $\Delta_{0,4}$.

These push forwards are easy to compute. A representative example will be given. Consider the graph R_4 :



with a fixed labelling of the marked points and the round markings. The three components of the graph have been labeled U , V , and W . The labeled graph now corresponds to an irreducible stratum R of X_ζ . An admissible cover lying above this stratum consists of the following data: an elliptic double cover of V ramified at a , b , c , and the intersection $V \cap U$, a rational double cover of U ramified at d and the intersection $V \cap W$, and an disjoint étale double cover of W . The double cover of W is a union of two $\mathbf{P}^1$'s. The markings $\{1, 2, 3\}$ can be distributed in 4 ways (up to isomorphism) on the 2 components over W :

$$(12, 3), (13, 2), (23, 1), (123).$$

This yields four components of $\pi^{-1}(R)$. The component $(12, 3)$ pushes forward by λ to:

$$6 \quad \begin{array}{c} | \quad 1 \ 2 \\ | \quad | \quad | \\ 3- \\ | \quad | \quad | \\ | \quad 4 \end{array} \quad 1$$

The λ push-forward of the component (123) lies in the divisor stratum of $\overline{M}_{1,4}$:

$$\begin{array}{c} | \\ | \quad 1 \\ | \quad 2 \\ | \quad 3 \\ | \\ | \quad | \quad | \\ | \quad 4 \end{array} \quad 1$$

and is easily seen to equal:

$$2 \quad \begin{array}{c} | \quad 1 \ 2 \\ | \quad | \quad | \\ 3- \\ | \quad | \quad | \\ | \quad 4 \end{array} \quad 1 \quad + \quad 2 \quad \begin{array}{c} | \quad 1 \ 3 \\ | \quad | \quad | \\ 2- \\ | \quad | \quad | \\ | \quad 4 \end{array} \quad 1 \quad + \quad 2 \quad \begin{array}{c} | \quad 2 \ 3 \\ | \quad | \quad | \\ 1- \\ | \quad | \quad | \\ | \quad 4 \end{array} \quad 1$$

Adding all the permutations and multiplying by the automorphism factor 2 associated to R_4 yields $\lambda_*\pi^*(R_4) = 16 \triangle_{2,3}$.

5 Getzler's relation

The calculations of $\lambda_*\pi^*$ in Section 4 and the linear equivalence (3) yield the relation:

$$48 \triangle_{2,2} - 16 \triangle_{2,3} - 4 \triangle_{2,4} + 12 \triangle_{3,4} \quad (4)$$

$$+ 2 \triangle_{0,2} + 10 \triangle_{0,3} + 8 \triangle_{0,4} - 6 \triangle_a - 16 \triangle_b = 0$$

in $A_2(\overline{M}_{1,4})$. The basic linear equivalence in $A_2(\overline{M}_{1,4})$ obtained from $\triangle_0$ is (see [G2]):

$$\triangle_{0,2} + 3 \triangle_{0,3} + 3 \triangle_{0,4} - 3 \triangle_a - 4 \triangle_b = 0 \quad (5)$$

Equation (4) minus twice equation (5) yields:

$$4 \cdot (12 \triangle_{2,2} - 4 \triangle_{2,3} - \triangle_{2,4} + 3 \triangle_{3,4} + \triangle_{0,3} + \frac{1}{2} \triangle_{0,4} - 2 \triangle_b) = 0$$

which is Getzler's relation [G2].

Theorem 1 *Getzler's relation in $H^4(\overline{M}_{1,4}, \mathbb{Q})$ is obtained from a rational equivalence in $A_2(\overline{M}_{1,4}, \mathbb{Q})$.*

6 Elliptic invariants of $\mathbf{P}^2$

For $d \geq 1$, let $N_d^{(0)}$ and $N_d^{(1)}$ be the rational and elliptic Gromov-Witten invariants of $\mathbf{P}^2$. Let Γ and E be the (quantum) rational and elliptic potentials:

$$\Gamma(y_1, y_2) = \sum_{d \geq 1} N_d^{(0)} e^{dy_1} \frac{y_2^{3d-1}}{(3d-1)!}, \quad (6)$$

$$E(y_1, y_2) = -\frac{y_1}{8} + \sum_{d \geq 3} N_d^{(1)} e^{dy_1} \frac{y_2^{3d}}{(3d)!}. \quad (7)$$

The variables y_0, y_1, y_2 correspond to the fundamental, line, and point classes of $\mathbf{P}^2$ respectively. The elliptic sum starts in degree 3 since $N_1^{(1)}$ and $N_2^{(1)}$ vanish. Let $\tilde{E} = E + \frac{1}{8}y_1$. The composition axiom for Gromov-Witten invariants ([KM], [RT], [BM]) and Getzler's relation immediately yield partial differential equations satisfied by the elliptic potential. Getzler uses these to obtain a complete recursion determining the numbers $N_d^{(1)}$ [G2]. The numbers $N_d^{(1)}$ have been obtained via more classical techniques in algebraic geometry in [CH], [R].

The relation predicted by Eguchi, Hori, and Xiong from the Virasoro conjecture is:

$$\frac{N_d^{(1)}}{(3d-1)!} = \frac{1}{12} \binom{d}{3} \frac{N_d^{(0)}}{(3d-1)!} + \sum_{d_1+d_2=d, d_i > 0} \frac{3d_1^2 d_2 - 2d_1 d_2}{9} \frac{N_{d_1}^{(0)}}{(3d_1-1)!} \frac{N_{d_2}^{(1)}}{(3d_2)!}. \quad (8)$$

While the equation suggests divisor geometry, no proof via such an approach is known to the author. The relation is established here via Getzler's equation and the WDVV-equations.

The first step is to rewrite the recursion (8) in differential form:

$$\tilde{E}_2 = \frac{\Gamma_{111} - 3\Gamma_{11} + 2\Gamma_1}{72} + \frac{\Gamma_{11}\tilde{E}_1}{3} - \frac{2\Gamma_1\tilde{E}_1}{9}.$$

The subscripts denote partial differentiation by y_1 and y_2 respectively. Next, the equation

$$\tilde{E}_1 = \frac{y_2}{3} \tilde{E}_2$$

is used to obtain:

$$\tilde{E}_2 \left(1 - \frac{1}{9}y_2\Gamma_{11} + \frac{2}{27}y_2\Gamma_1\right) = \frac{\Gamma_{111} - 3\Gamma_{11} + 2\Gamma_1}{72}.$$

Finally, we see:

$$E = -\frac{y_1}{8} + \int \frac{\Gamma_{111} - 3\Gamma_{11} + 2\Gamma_1}{72} \left(1 - \frac{1}{9}y_2\Gamma_{11} + \frac{2}{27}y_2\Gamma_1\right)^{-1} dy_2. \quad (9)$$

Formula (9) is equivalent to (8).

In order to prove (8), we consider a differential equation obtained from Getzler's relation. Let

$$\pi : \overline{M}_{1,4+3d-6}(\mathbf{P}^2, d) \rightarrow \overline{M}_{1,4}$$

be the natural projection. Let

$$e_i : \overline{M}_{1,4+3d-6} \rightarrow \mathbf{P}^2$$

be the natural evaluation maps. The codimension 2 Gromov-Witten class

$$\pi_* \left([\overline{M}_{1,4+3d-6}(\mathbf{P}^2, d)]^{Vir} \cap \prod_{i=1}^{3d-2} e_i^*([\text{point}]) \right)$$

intersected with Getzler's relation yields the following differential equation:

$$\begin{aligned} & 36E_{11}\Gamma_{122}^2 - 48E_{12}\Gamma_{112}\Gamma_{122} - 48E_{22}\Gamma_{222} - 12E_1\Gamma_{1122}\Gamma_{122} \\ & + 24E_1\Gamma_{112}\Gamma_{1222} + 24E_2\Gamma_{2222} + 2\Gamma_{1222}\Gamma_{1112} \\ & + \frac{1}{2}\Gamma_{12222}\Gamma_{111} + \frac{3}{2}\Gamma_{22222} - 3\Gamma_{1122}^2 = 0 \end{aligned} \quad (10)$$

It is easily seen by examining the coefficient recursions that (10) uniquely determines the potential E in form (7) from Γ . Therefore, to establish (8), it suffices to prove (9) determines a solution of (10).

Replace the terms E_1 , E_{11} , E_{12} in (10) by:

$$\begin{aligned} E_1 &= \frac{y_2}{3}E_2 - \frac{1}{8}, \\ E_{11} &= \frac{y_2^2}{9}E_{22} + \frac{y_2}{9}E_2, \\ E_{12} &= \frac{y_2}{3}E_{22} + \frac{1}{3}E_2. \end{aligned}$$

Equation (9) then may be used to replace the terms E_2 and E_{22} by derivatives of Γ . A partial differential equation for Γ is then obtained. Equation (9) defines a solution of (10) if and only if Γ satisfies this differential equation.

The WDVV-equation for Γ is:

$$\Gamma_{222} - \Gamma_{112}^2 + \Gamma_{111}\Gamma_{122} = 0. \quad (11)$$

Γ also satisfies:

$$\Gamma_1 = \frac{y_2}{3}\Gamma_2 + \frac{1}{3}\Gamma. \quad (12)$$

A simple check in symbolic algebra shows the required differential equation for Γ is implied by the WDVV-equation and its first two y_2 -derivatives together

with (12). In fact, the required differential equation for Γ modulo equation (12) is seen to be of the form:

$$A\Psi + B\Psi_2 + C\Psi_2^2 + D\Psi_{22} = 0. \quad (13)$$

A , B , C , and D are polynomials in y_2 , Γ , and the derivatives of Γ . Ψ is the left side of the WDVV-relation (11), and the subscripts denote partial differentiation by y_2 .


$$\begin{aligned} A &= -2^7 \cdot (36y_2\Gamma_{22} + 24y_2^2\Gamma_{222} - 4\Gamma_1\Gamma_{11} - 9\Gamma_{11}^2 + \frac{4}{3}\Gamma_1^2 \\ &\quad + 12\Gamma_1\Gamma_{111} - 18\Gamma_{11}\Gamma_{111} + 27\Gamma_{111}^2) \\ B &= -2^6 \cdot (-60\Gamma + 375\Gamma_1 - \frac{1521}{2}\Gamma_{11} + \frac{783}{2}\Gamma_{111} \\ &\quad + 2y_2\Gamma_1^2 - 6y_2\Gamma_1\Gamma_{11} + 8y_2\Gamma_1\Gamma_{111} \\ &\quad - \frac{11}{2}y_2\Gamma_{11}^2 - 9y_2\Gamma_{11}\Gamma_{111} + \frac{9}{2}y_2\Gamma_{111}^2) \\ C &= -2^6 \cdot y_2^4 \\ D &= -2^5 \cdot (y_2\Gamma_{111} + 9) \cdot (3y_2\Gamma_{11} - 2y_2\Gamma_1 - 27) \end{aligned}$$


This concludes the proof of recursion (8).

Theorem 2 *The Eguchi-Hori-Xiong formula holds for the elliptic Gromov-Witten invariants of $\mathbf{P}^2$.*

Appendix A Strata of $\overline{M}_{1,4}$ in dimension 2

Getzler's notation for the complete set of $\mathbb{S}_4$ -invariant dimension 2 strata of $\overline{M}_{1,4}$ is followed:





The short bars indicate the marked points $\{1, 2, 3, 4\}$. Nonsingular genus 1 components (labeled by 1) occur in the first row. Each graph denotes the sum of strata in the corresponding $\mathbb{S}_4$ -orbit. For example, $\Delta_{2,2}$ and $\Delta_{2,3}$ are sums of 3 and 12 strata respectively.

Appendix B Strata of X_ζ in dimension 2


A graph G below *together* with an assignment μ of the round markings to the set $\{a, b, c, d\}$ corresponds to an $\mathbb{S}_4$ -invariant dimension 3 stratum of $\overline{M}_{0,8}$. The 8 point marking set is $\{1, 2, 3, 4, a, b, c, d\}$ as in Section 1, and the $\mathbb{S}_4$ -action is on the first four markings. By intersecting these strata of $\overline{M}_{0,8}$ with X_ζ , a sum of dimension 2 strata of X_ζ is associated to (G, μ) .


For example, consider R_3 . After an assignment of the round markings to the set $\{a, b, c, d\}$, there are 4 strata of $\overline{M}_{0,8}$ in the $\mathbb{S}_4$ -orbit. These yield a sum of four strata of X_ζ . Similarly, S_3 together with an assignment of the round markings yields a sum 12 strata of X_ζ .

The assignment μ of the round markings will be suppressed since the calculation of the map

$$\lambda_* \pi^* : A_2(X_\zeta) \rightarrow A_2(\overline{M}_{1,4})$$

does not depend upon μ . Let $K = \text{Ker}(\lambda_* \pi^*)$. Let μ and μ' be two round marking assignments for a fixed graph G . The two cycles corresponding to (G, μ) and (G, μ') are equal in $A_2(X_\zeta)/K$.






This is not a complete list. Only the strata which are required for the computations in Section 3 are given.

Appendix C Strata of X_ζ in dimension 3

As in Appendix B, a graph below together with an assignment of the round markings to the set $\{a, b, c, d\}$ corresponds to a sum of dimension 3 strata of X_ζ .



References

- [CH] L. Caporaso and J. Harris, *Counting plane curves of any genus*, preprint 1996.
- [BM] K. Behrend and Yu. Manin, *Stacks of stable maps and Gromov-Witten invariants*, Duke J. Math. **85** (1996), 1-60.

- [EHX] T. Eguchi, K. Hori, and C.-S. Xiong, *Quantum cohomology and Virasoro algebra*, preprint 1997.
- [Fa] C. Faber, *Chow rings of moduli spaces of curves. I. The Chow ring of $\overline{M}_3$* , Ann. of Math. **132** no. 2 (1990), 331-419.
- [Fu] W. Fulton, *Intersection theory*, Springer-Verlag: Berlin, 1984.
- [FM] W. Fulton and R. MacPherson, *A compactification of configuration spaces*, Ann. of Math. **139** no. 1 (1994), 183-225.
- [FP] W. Fulton and R. Pandharipande, *Notes on stable maps and quantum cohomology*, Proc. AMS (to appear).
- [G1] E. Getzler, *The semi-classical approximation for modular operads*, preprint 1996.
- [G2] E. Getzler, *Intersection theory on $\overline{M}_{1,4}$ and elliptic Gromov-Witten invariants*, preprint 1996.
- [HM] J. Harris and D. Mumford, *On the Kodaira dimension of the moduli space of curves*, Invent. Math. **67** (1982), 23-86.
- [K] M. Kontsevich, *Enumeration of rational curves via torus actions*, in *The moduli space of curves*, R. Dijkgraaf, C. Faber, and G. van der Geer, eds., Birkhauser, 1995, pp 335-368.
- [KM] M. Kontsevich and Yu. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Commun. Math. Phys. **164** (1994), 525-562.
- [R] Z. Ran, *The degree of a Severi variety*, Bull. Amer. Math. Soc. **17** no. 1 (1987), 125-128.
- [RT] Y. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, J. Diff. Geom. **42** (1995), 259-367.
- [V] A. Vistoli, *Chow groups of quotient varieties*, J. Algebra **107** no. 2 (1987), 410-424.

Department of Mathematics
University of Chicago
5734 S. University Ave. 60637
rahul@math.uchicago.edu